

LETTER • OPEN ACCESS

Daytime cooling efficiencies of urban trees derived from land surface temperature are much higher than those for air temperature

To cite this article: Meng Du *et al* 2024 *Environ. Res. Lett.* **19** 044037

View the [article online](#) for updates and enhancements.

You may also like

- [Trees in cool climate cities may increase atmospheric carbon by altering building energy use](#)
Tedward Erker and Philip A Townsend
- [Global assessment of urban trees' cooling efficiency based on satellite observations](#)
Qiquan Yang, Xin Huang, Xiaohua Tong et al.
- [The use of mobile lidar data and Gaofen-2 image to classify roadside trees](#)
Minye Wang, Rufe Liu, Xiushan Lu et al.

The Breath Biopsy® Guide
Fourth edition

FREE

DOWNLOAD THE FREE E-BOOK

BREATH BIOPSY

OWLSTONE MEDICAL

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

OPEN ACCESS

RECEIVED
17 November 2023REVISED
22 February 2024ACCEPTED FOR PUBLICATION
6 March 2024PUBLISHED
22 March 2024

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Daytime cooling efficiencies of urban trees derived from land surface temperature are much higher than those for air temperature

Meng Du¹, Niantan Li¹, Ting Hu¹, Qiquan Yang^{2,3,4,*} , TC Chakraborty⁵, Zander Venter⁶  and Rui Yao⁷¹ School of Remote Sensing and Geomatics Engineering, Nanjing University of Information Science and Technology, Nanjing 210044, People's Republic of China² College of Surveying & Geo-Informatics, Tongji University, Shanghai 200092, People's Republic of China³ State Key Laboratory of Lunar and Planetary Sciences, Macau University of Science and Technology, Macau, People's Republic of China⁴ The Shanghai Key Laboratory of Space Mapping and Remote Sensing for Planetary Exploration, Tongji University, Shanghai 200092, People's Republic of China⁵ Pacific Northwest National Laboratory, Richland, WA, United States of America⁶ Norwegian Institute for Nature Research, Trondheim, Norway⁷ School of Remote Sensing and Information Engineering, Wuhan University, Wuhan 430079, People's Republic of China

* Author to whom any correspondence should be addressed.

E-mail: qiqquanyang@tongji.edu.cn**Keywords:** urban heat islands, green space, *in situ* measurements, ecosystem service, mitigation strategies, remote sensing**Abstract**

Accurately capturing the impact of urban trees on temperature can help optimize urban heat mitigation strategies. Recently, there has been widespread use of remotely sensed land surface temperature (T_s) to quantify the cooling efficiency (CE) of urban trees. However, remotely sensed T_s reflects emitted radiation from the surface of an object seen from the point of view of the thermal sensor, which is not a good proxy for the air temperature (T_a) perceived by humans. The extent to which the CEs derived from T_s reflect the true experiences of urban residents is debatable. Therefore, this study systematically compared the T_s -based CE (CE_{T_s}) with the T_a -based CE (CE_{T_a}) in 392 European urban clusters. CE_{T_s} and CE_{T_a} were defined as the reductions in T_s and T_a , respectively, for every 1% increase in fractional tree cover (FTC). The results show that the increase in FTC has a substantial impact on reducing T_s and T_a in most cities during daytime. However, at night, the response of T_s and T_a to increased FTC appears to be much weaker and ambiguous. On average, for European cities, daytime CE_{T_s} reaches $0.075\text{ }^\circ\text{C}\ \%^{-1}$, which is significantly higher (by an order of magnitude) than the corresponding CE_{T_a} of $0.006\text{ }^\circ\text{C}\ \%^{-1}$. In contrast, the average nighttime CE_{T_s} and CE_{T_a} for European cities are similar, both approximating zero. Overall, urban trees can lower daytime temperatures, but the magnitude of their cooling effect is notably amplified when using remotely sensed T_s estimates compared to *in situ* T_a measurements, which is important to consider for accurately constraining public health benefits. Our findings provide critical insights into the realistic efficiencies of alleviating urban heat through tree planting.

1. Introduction

The process of urbanization, marked by changes in land cover and the release of anthropogenic heat emissions, typically results in elevated temperatures within urban areas, giving rise to the phenomenon known as the urban heat island (UHI) effect (Kalnay and Cai 2003, Grimm *et al* 2008, Zhang *et al* 2022b). This additional urban warming can have negative impacts on various aspects of urban environments, including the vegetation growth, the hydrological

cycle, and the soil quality (Zhou *et al* 2018, Gui *et al* 2019, Zou *et al* 2021). Moreover, urban heat can pose a risk to the well-being of urban residents, influencing their indoor and outdoor thermal comfort and even leading to added mortality (Zhou *et al* 2018, Iungman *et al* 2023, Ren *et al* 2023). Therefore, urban heat and strategies for its mitigation have garnered increased attention in recent years (Yao *et al* 2018, Zhou *et al* 2018, Hsu *et al* 2021, Tuholske *et al* 2021, Chakraborty *et al* 2022c, Liu *et al* 2023, Schneider *et al* 2023, Yang *et al* 2023). Increasing vegetation

cover, particularly through tree planting, is a feasible and commonly proposed approach for alleviating urban heat (Schwaab *et al* 2021, Lungman *et al* 2023, Zhang *et al* 2023). Trees outperform other vegetation in terms of evapotranspiration and shading, making them effective in reducing local temperatures (Zhou *et al* 2017, Liu *et al* 2021, Zhang *et al* 2023). Consequently, the study of the cooling effect (CE) of trees is a prominent and actively researched topic in the field of urban climate.

Traditionally, the cooling efficiency of trees is quantified by directly comparing the air temperature (T_a) between areas with and without trees using *in situ* measurements (Akbari *et al* 1990, Richards *et al* 2020, Yan *et al* 2020, Zhang *et al* 2020). Such approach offers the advantage of continuous, long-term observations and yields results that align closely with human perception (Goldblatt *et al* 2021). However, *in situ* measurements of the thermal environment (e.g. weather stations or mobile transects) are usually characterized by spatial discontinuities and limited coverage, making them too costly for extensive large-scale analyses (Peters *et al* 2010, Loughner *et al* 2012, Wang *et al* 2012, 2018, Jiao *et al* 2017, Richards *et al* 2020, Wujeska-Klaue and Pfautsch 2020, Yan *et al* 2020, Zhang *et al* 2020, Meili *et al* 2021). With the proliferation of satellite observations, remotely sensed land surface temperature (T_s) has been often used for quantifying the CE of urban trees due to its advantages of spatial continuity (Wang *et al* 2019, 2020, Yang *et al* 2022). Nevertheless, it is crucial to acknowledge that a distinction exists between remotely sensed T_s observations and *in situ* T_a measurements (Jin and Dickinson 2010, Jin 2012, Zhang *et al* 2014, Chakraborty *et al* 2017, Janatian *et al* 2017, Goldblatt *et al* 2021, Venter *et al* 2021, Naserikia *et al* 2023).

The differences between T_s and T_a are primarily evident in two aspects. Firstly, T_s and T_a vary in magnitude and spatial distributions, originating from factors like radiation, heat transfer, and storage processes (Mutiibwa *et al* 2015, Cao *et al* 2021). Solar radiation absorbed by surface materials with high emissivity results in a significant increase in surface temperature, whereas air, with its lower emissivity, is less efficient at absorbing and radiating energy (Harde 2013, Li *et al* 2013). Heat transfer via conduction and convection contributes to temperature interaction, but the efficiency of this process is limited by air's low thermal conductivity (Zhou *et al* 2012, Lu *et al* 2014). The higher thermal capacity of surface materials enables them to absorb more heat during the day compared to air, thereby amplifying temperature discrepancies between the surface and the surrounding air (Mohajerani *et al* 2017). Secondly, T_s and T_a represent different components of the environment, with distinct impacts on human comfort. T_s is typically not directly experienced by urban pedestrians since we do not come into direct contact with the surfaces using our bare skin (Goldblatt *et al* 2021, Haeri *et al*

2023). Moreover, T_s measurements often relate to elevated locations like rooftops, walls, and the top of tree canopies, areas not frequented by humans for daily activities (Vanos *et al* 2016, Battisti *et al* 2018, Haeri *et al* 2023). As a result, the CE of trees derived through T_s may not appropriately capture the public health impacts on urban inhabitants, which are typically more aligned with the T_a perceived by humans (Novick and Katul 2020, Chakraborty *et al* 2022c).

Although multiple numerical modeling studies have estimated the CE of urban trees (Krayenhoff *et al* 2021), observational estimates are much rarer (Ziter *et al* 2019, Chakraborty *et al* 2022c), partly due to the dearth of dense T_a measurements within cities (Zaitchik and Tuholske 2021). As a result, many researchers have quantified the CE of urban trees using remotely sensed T_s observations and considered the slope of T_s versus tree cover fraction to quantify CE (Wang *et al* 2019, 2020, Chinchilla *et al* 2021, Schwaab *et al* 2021, Zhou *et al* 2021, Chakraborty *et al* 2022a, Zhao *et al* 2023a). Since T_s and T_a are different, we would also expect substantial differences in T_a -based CE (CE_{T_a}) and T_s -based CE (CE_{T_s}). Knowledge about how CE_{T_s} deviates from CE_{T_a} is critical for gaining a more physiologically-relevant understanding of the role played by trees in shaping the urban thermal environment. However, there is still a lack of large-scale and quantitative assessments of the relationship and difference between CE_{T_s} and CE_{T_a} . The study by Chakraborty *et al* (2022a) is the only relevant one, but it utilizes a vegetation index instead of tree cover fraction, with the two being not directly comparable (Zeng *et al* 2023).

Hence, by using spatiotemporally matched *in situ* T_a and remotely sensed T_s datasets, we calculated CE_{T_a} and CE_{T_s} , respectively, in 392 European urban clusters. Subsequently, we conducted a comparative analysis of the differences between CE_{T_a} and CE_{T_s} in terms of spatial distribution and day-night contrast. The results of this study can deepen our understanding of the cooling benefits of urban trees and help inform heat stress mitigation strategies.

2. Data and methods

2.1. Data selection and processing

The fractional tree cover (FTC) was obtained from the Copernicus Global Land Service (CGLS) dataset (2019). Based on over 20 000 random samples, Buchhorn *et al* (2020) reported that the CGLS dataset had an overall accuracy of better than 80% for all land-cover types and a global average absolute error of 9% for the FTC (Buchhorn *et al* 2020). The surface water was derived from the global surface water data produced by the Joint Research Centre (Pekel *et al* 2016), which provides annual maximum surface water extents. Surface elevation information




Figure 1. Spatial patterns and frequency distributions of the daytime cooling efficiency (CE) of urban trees across European urban clusters. CE_{T_s} and CE_{T_a} represent the CE estimated from surface temperature and the CE estimated from air temperature, respectively. The observational times are $\sim 13:30$ for AQUA and $\sim 10:30$ for TERRA, respectively. The air temperature corresponds to overpassing time of the overlaying MODIS pixels.

was obtained from Global 30 Arc-Second Elevation (GTOPO30) dataset.

The growing popularity of Citizen Weather Stations (CWSs) in urban areas in recent years has facilitated the crowdsourcing of intensive meteorological observations. Venter *et al* (2021) obtained hourly T_a data for July 2019 from all available CWS provided by Netatmo in Europe (Chakraborty *et al* 2022b). The CWS T_a data underwent additional cleaning processing to remove stations with poor data quality or significant missing data (Venter *et al* 2021). This led to the final selection of 75 293 CWS stations, distributed across 931 European urban clusters. The boundaries of urban clusters were derived from the Global Human Settlement Layer dataset using the clustering method proposed by Venter *et al* (2021).

Daily T_s data for the European urban clusters during July 2019 were collected from the MODIS products (MOD11A1 and MYD11A1). These T_s

data were preprocessed into clear-sky pixels with an average T_s error less than or equal to 3 K. MODIS satellites collected data every day around 1:30, 10:30, 13:30, and 22:30 local time to provide daytime and nighttime T_s observations. To obtain the hour-by-hour T_s values, the MODIS transit time from local solar time was converted to UTC local time by subtracting the longitude of the MODIS grid and dividing by 15, according to the MODIS user guide (Wan 2006). The time zones of each urban cluster were then accounted for and combined with daylight saving time to offset the UTC of each city. This resulted in T_s readings with specific hourly timestamps that could be synchronized with the hourly CWS dataset (Venter *et al* 2021, Chakraborty *et al* 2022b).

The T_s pixels were paired with the treated CWS T_a stations based on coordinates and time (Chakraborty *et al* 2022c). Subsequently, we calculated the median values of T_s and T_a at each




Figure 2. Spatial patterns and frequency distributions of the nighttime cooling efficiency (CE) of urban trees across European urban clusters. CE_{T_s} and CE_{T_a} represent the CE estimated from surface temperature and the CE estimated from air temperature, respectively. The observational times are $\sim 1:30$ for AQUA and $\sim 22:30$ for TERRA, respectively. The air temperature corresponds to overpassing time of the overlaying MODIS pixels.

station for the study period separately. We chose the median rather than the mean to avoid the effects of extreme temperatures. Then, we extracted information about tree cover fraction, elevation, and the presence of surface water for each station. In each urban clusters, we removed stations covered by surface water or influenced by topographic relief (out of the range of mean elevation ± 100 m) (Cao *et al* 2016, Li *et al* 2022). In addition, we removed urban clusters with too few stations (< 20) to ensure the stability of the regression model for calculating the CE of urban trees. The above processes reduced the number of available stations to 40 364. Ultimately, a total of 392 urban clusters were included in this study (figure 1). It should be emphasized that the MOD11A1 T_s had data gaps during nighttime due to cloud contamination and quality degradation, resulting in an insufficient number of observations (< 20) in some urban clusters. Therefore, we calculated the CE cor-

responding to the MOD11A1 nighttime T_s ($\sim 22:30$) only for the 232 urban clusters (figure 2).

2.2. Calculation of cooling efficiency of urban trees

In this study, we employed a univariate linear regression model to calculate the CE_{T_s} and CE_{T_a} for each urban cluster. In this model, we considered FTC as the independent variable, and either T_s or T_a as the dependent variable. We computed the regression coefficients and used the opposite of these coefficients as the CE of urban trees, denoted as $CE_{T_s} = -\Delta T_s / \Delta \text{FTC}$ and $CE_{T_a} = -\Delta T_a / \Delta \text{FTC}$. CE_{T_s} and CE_{T_a} represent the reduction in T_s and T_a , respectively, for each 1% increase in the percentage of urban trees, and measure the CE of urban trees with respect to T_s and T_a (Wang *et al* 2019, 2020, Cheng *et al* 2022, Yang *et al* 2022). If CE_{T_s} is higher than CE_{T_a} , it suggests a higher estimated CE of urban trees based on T_s compared to T_a , and vice versa.




Figure 3. Comparisons of the cooling efficiencies of urban trees estimated from surface temperature (CE_{T_s}) with that estimated from air temperature (CE_{T_a}). The p -values are from significance t-tests between CE_{T_s} and CE_{T_a} . If the p -value is less than 0.05, it is considered a statistically significant difference between CE_{T_s} and CE_{T_a} . The lines and dots in the boxes represent median and mean values of trees' cooling efficiency, respectively. The error bars indicate 95% confidence intervals. The observational times are $\sim 13:30$ and $\sim 1:30$ for AQUA, and $\sim 10:30$ and $\sim 22:30$ for TERRA. The air temperature measurements corresponding to overlaying MODIS pixels are used for daytime and nighttime overpasses.

3. Results

During daytime, both CE_{T_s} and CE_{T_a} show positive values in most urban clusters, indicating the prevailing cooling effect exerted by urban trees on local temperatures (figure 1). Statistically, over 90% urban clusters exhibit positive daytime CE_{T_s} , whereas the corresponding percentage for daytime CE_{T_a} stands at approximately 60% (figure 1). Besides, higher values of CE_{T_s} and CE_{T_a} are mostly concentrated in western Europe and south-central Europe. On average for all urban clusters, CE_{T_s} reaches 0.075 [$0.069, 0.082$] $^{\circ}\text{C } \%^{-1}$ (95% confidence intervals in parentheses, hereafter) and 0.055 [$0.049, 0.061$] $^{\circ}\text{C } \%^{-1}$ at local times $\sim 13:30$ and $\sim 10:30$, respectively (figure 3). These mean values of CE_{T_s} are significantly ($p < 0.001$, t-test) higher than corresponding averages of CE_{T_a} , which are 0.006 [$0.001, 0.010$] $^{\circ}\text{C } \%^{-1}$ and 0.003 [$-0.001, 0.008$] $^{\circ}\text{C } \%^{-1}$, respectively (figure 3). These results indicate that the daytime CE of urban trees quantified from T_s differs from that estimated from T_a , which is further supported by the weak correlation between CE_{T_s} and CE_{T_a} across all urban clusters (figure 4).

At night, CE_{T_a} is found to be positive in over 70% urban clusters, while CE_{T_s} does not exhibit a fixed positive or negative tendency (figure 2). In addition, higher values of CE_{T_s} and CE_{T_a} are concentrated in western Europe and south-central Europe. On average for all urban clusters, CE_{T_s} reaches 0.018 [$0.015, 0.022$] $^{\circ}\text{C } \%^{-1}$ and -0.004 [$-0.007, -0.002$] $^{\circ}\text{C } \%^{-1}$ at local times $\sim 13:30$ and $\sim 10:30$, respectively (figure 3). The mean values of these CE_{T_s} did not differ significantly from the corresponding averages for CE_{T_a} , which are 0.018 [$0.014, 0.021$] $^{\circ}\text{C } \%^{-1}$ and 0.008 [$0.005, 0.011$] $^{\circ}\text{C } \%^{-1}$, respectively (figure 3). These


results indicate that the nighttime CE of urban trees quantified using T_s has a higher similarity to that estimated from T_a compared to the daytime results. This finding is further supported by the observation that the distribution of CE_{T_s} and CE_{T_a} aligns more closely with the 1:1 line during nighttime than during daytime (figure 4).

To provide a more intuitive representation of the difference between CE_{T_s} and CE_{T_a} , we selected six prototypical urban clusters (Berlin, Paris, Reims, Budapest, Bratislava, and Wien) with different climates, tree species, and city sizes for closer examination (figure 5). It can be seen that both daytime T_s and T_a show almost a consistent decreasing trend with increasing FTC, suggesting that trees can play a role in reducing urban localized temperatures. However, T_s typically exhibits a greater range of spatial variability relative to T_a , and the extent of temperature change triggered by an equivalent percentage increase in FTC is more pronounced for T_s compared to T_a . This discrepancy in the response of T_s and T_a to increased FTC ultimately leads to the observed difference between CE_{T_s} and CE_{T_a} .

4. Discussion


4.1. Spatiotemporal difference in the cooling efficiencies estimated from T_s and T_a

The findings suggest a noticeable CE of urban trees during daytime, and the impact of trees on reducing T_s is found to be substantially more than that on T_a . Trees release water vapor through small pores called stomata in their leaves, also known as transpiration (Pirasteh-Anosheh et al 2016). As this water evaporates, it absorbs heat from the surroundings, lowering the temperature of the surface, which in



turn impacts the near surface (Gunawardena *et al* 2017). Evapotranspiration occurs in the immediate vicinity of the ground, where trees draw up water from the soil and release water vapor through stomata in their leaves (Berry *et al* 2010). This close proximity to the surface means that the CE is more directly felt at ground level (Armson *et al* 2012). While water vapor released through evapotranspiration does influence T_a , the impact is somewhat diluted as it mixes with the larger volume of the atmosphere (Gunawardena *et al* 2017). Therefore, the CE for T_a is not as concentrated as it is at the surface. Second, trees provide shade by blocking sunlight from reaching the ground, which reduces the amount of solar radiation absorbed by the Earth's surface, leading to a substantial CE (Coutts *et al* 2016). On the other hand, T_a is influenced by a variety of factors, including the absorption of solar radiation by the atmosphere, conduction, convection, and radiation (Raman *et al* 2014). While shading by trees can have

some influence on the T_a by reducing the amount of solar radiation reaching the ground, the atmosphere itself is less directly affected by shading compared to the surface (Rahman *et al* 2017). Therefore, the shade provided by trees has a more direct impact on the T_s than on the T_a . However, we should note here that the T_s underneath the tree can be markedly different from the T_s measured by satellites, whose view of the ground can be partially blocked by those trees. Third, trees play a pivotal role in creating microclimates by regulating soil moisture in their immediate surroundings (Pramova *et al* 2012). Adequate soil moisture contributes to a cooler surface, and the impact of soil moisture regulation is particularly significant at the surface level, where trees interact with the ground (Lagergren and Lindroth 2002). In summary, the creation of microclimates by trees involves a combination of factors, including the evapotranspiration process, soil moisture control, and shade provision. While these factors also influence T_a , their immediate



and more pronounced effects are often observed at the surface level where the vegetation interacts intimately with the ground.

However, the CE of urban trees on nighttime temperatures is very limited in most cities. During the night, in the absence of sunlight for photosynthesis, trees generally close their stomata (Domec *et al* 2012). As a consequence of stomatal closure, metabolic activities in the leaves, including transpiration, slow down or cease (Dawson *et al* 2007). This reduction in transpiration leads to a notable decrease in the CE facilitated by this process (Zheng *et al* 2021). In addition, trees, especially deciduous ones, typically have a lower albedo, which results in more energy absorption during the day (Hami *et al* 2019). The additional absorbed energy is then released at night, thereby offsetting the cooling benefits that trees typically provide (Peng *et al* 2014, Li *et al* 2015). Furthermore, in urban environments, tall and dense trees can influence the dissipation of heat from various structures. The obstruction of airflow may contribute to heat retention in certain areas, especially during calm nights (Cleugh 1998, Zhao *et al* 2023b). The factors mentioned above, to some extent, offer a plausible explanation for the reduced CE of trees

at night. This also underscores the complex interplay of various factors in shaping the overall thermal behavior of urban environments influenced by tree cover.

We have identified notable variations in the CE of urban trees across different cities. Firstly, the daytime CE is found to be higher in western Europe compared to central Europe (figure 1). This difference may be attributed to variations in tree species. Western Europe is dominated by oaks from broadleaf forests (<https://inventaire-forestier.ign.fr/spip.php?rubrique86>), while central Europe is dominated by spruce and pine from coniferous forests (Holzwarth *et al* 2020). Characteristics of urban trees, such as leaf structure, color, size, and density, directly influence their effects on the surrounding environment and the observed CE (Liu *et al* 2019, Wang *et al* 2021). Secondly, a relatively higher CE is also observed in south-central Europe, characterized by a hotter and drier environment. This may be attributed to the nonlinear physiological response of urban trees to meteorological conditions, which generally manifests itself as an increase in evapotranspiration rates with increasing temperature and decreasing humidity (Wang *et al* 2019, 2020, Yang *et al* 2022).

Besides, the role of urban trees in moderating temperature is also influenced by their spatial configurations, with densely and continuously distributed trees tending to provide more cooling benefits (Zhou et al 2017, Fan et al 2019). This may provide a plausible explanation for the significantly higher CE in some well-greened cities, such as Ludenscheid and Sodertalje (figure 1).

4.2. Significance and implications

This study finds that the estimated daytime CE of urban trees, based on T_a , is much lower than that based on T_s , which is in line with field-level estimates for general afforestation (Novick and Katul 2020) and multi-city estimates based on vegetation index (Chakraborty et al 2022c). Many recent multi-city studies have primarily utilized T_s instead of T_a to assess the cooling efficiency of trees (Wang et al 2019, 2020, Chinchilla et al 2021, Schwaab et al 2021, Zhou et al 2021, Cheng et al 2022, Chakraborty et al 2022a, Zhao et al 2023a). However, T_a is more closely related to temperatures perceived by humans than T_s . Hence, the current T_s -based CE might inflate the cooling effectiveness of tree planting and thus overestimate their public health benefits. This could potentially foster an excessively optimistic perspective regarding the cooling advantages associated with urban trees (Venter et al 2021, Chakraborty et al 2022c, Wang et al 2023a). It should be noted that increased FTC seems to be associated with a more efficient cooling of temperatures in western and south-central Europe. Therefore, emphasizing the increase of tree cover in these regions could be crucial for mitigating the adverse effects of high temperatures on human well-being. However, overall, it appears that the increase in tree cover in most parts of Europe is not significantly associated with a reduction in T_a . Particularly at night, there is even evidence of a warming effect of tree cover in some cities. Consequently, relying solely on increasing tree cover may not be sufficient for alleviating urban heat stress, and a more comprehensive approach may be necessary. Policymakers should consider implementing additional measures to effectively address urban heat. Feasible practical measures may involve enhancing surface albedo through the adoption of lighter roofing and paving materials, along with reducing human-generated heat emissions through promoting energy-saving lifestyles (Hayes et al 2022).

While we focus on urban vegetation impacts on T_a and T_s in the present study, there are other aspects of heat exposure within cities that should be kept in mind. Of note, urban trees can also increase humidity through transpiration (Meili et al 2020), which, in turn, can increase moist heat stress, all else remaining equal (Ho et al 2016, Chakraborty et al 2022c). While T_s should not directly be used as a proxy for heat stress, T_s has been successfully used as input to data-driven models to estimate urban T_a (Venter

et al 2020, Zhang et al 2022a) and even humidity (Wang et al 2023b). Trees can also provide shading, which during midday, can have a dominant impact on reducing overall heat loading (Middel et al 2021). However, T_a is only indirectly related to shading and satellites cannot sufficiently resolve shading from trees since they are primarily observing the top of the tree canopy. A recent study (Li et al 2023) over Philadelphia, USA estimated mean radiant temperature (T_{mrt}), a metric for thermal discomfort that also accounts for radiant heat, and found that satellite-derived T_s showed medium associations with T_{mrt} . A future direction may be to examine quantitative differences and similarities between the CE for T_s and T_{mrt} .

4.3. Limitations and future studies

The present study has certain limitations that should be kept in mind. First, it is important to note that the study area is limited to Europe, and therefore our findings cannot be generalized to other climate zones globally. Second, this study only considers data from the summer month of July, overlooking seasonal variations in vegetation activity, which are temperature-dependent. It is crucial to conduct in-depth investigations into the similarities and differences in CE quantified using T_s and T_a for other seasons to capture the full spectrum of potential variations. Third, there is a discrepancy in the spatial continuity and scale between satellite T_s observations and *in situ* T_a measurements, and a more detailed analysis considering variations at different spatial scales could provide additional insights. Fourth, there are sampling biases in the CWS data which are difficult to correct for. The most prominent bias is that CWS are typically placed in residential areas, rather than in commercial districts, which tend to experience higher temperatures (as indicated by T_s) (Hulley et al 2019, Chakraborty et al 2022c). This means that when utilizing CWS data, we might systematically overlook non-residential areas where pedestrians could still face higher-than-expected heat stress. To ensure the accurate utilization of CWS in urban research, it is essential to improve the metadata for the sensor locations. Furthermore, there is a pressing need for more robust quality control measures. To address the aforementioned limitations, future investigations should incorporate large-scale, multi-seasonal and multi-scale analyses from multiple data sources, to support more robust constraints on the physiologically-relevant cooling efficiency of urban trees.

Data availability statements

The data that support the findings of this study are openly available. The CWS air temperature measurements and corresponding MODIS land surface temperature observations are available at https://github.com/TC25/Europe_CWS. The CGLS fractional tree

cover data can be downloaded from <https://zenodo.org/records/3939050>. The JCR global surface water data can be accessed from <https://global-surface-water.appspot.com/download>. The Global 30 Arc-Second Elevation data is available from www.usgs.gov/centers/eros/science/usgs-eros-archive-digital-elevation-global-30-arc-second-elevation-gtopo30.

Acknowledgments

This research was supported by the Postdoctoral Science Foundation of China (2021TQ0245 and 2021M702470), and the National Natural Science Foundation of China (42201389 and 42201377). T.C.'s contribution was supported by the U.S. Department of Energy (DOE), Office of Science, Biological and Environmental Research program through the Early Career Research Program. Pacific Northwest National Laboratory is operated for DOE by Battelle Memorial Institute under contract DE-AC05-76RL01830.

ORCID iDs

Qiquan Yang  <https://orcid.org/0000-0003-1152-5999>

Zander Venter  <https://orcid.org/0000-0003-2638-7162>

References

- Akbari H, Rosenfeld A H and Taha H 1990 Summer heat islands, urban trees, and white surfaces (<https://doi.org/10.1103/physrevd.41.2057>)
- Armson D, Stringer P and Ennos A 2012 The effect of tree shade and grass on surface and globe temperatures in an urban area *Urban For. Urban Green.* **11** 245–55
- Battisti A, Laureti F, Zinzi M and Volpicelli G 2018 Climate mitigation and adaptation strategies for roofs and pavements: a case study at Sapienza University Campus *Sustainability* **10** 3788
- Berry J A, Beerling D J and Franks P J 2010 Stomata: key players in the earth system, past and present *Curr. Opin. Plant Biol.* **13** 232–9
- Buchhorn M, Lesiv M, Tsendbazar N-E, Herold M, Bertels L and Smets B 2020 Copernicus global land cover layers—collection 2 *Remote Sens.* **12** 1044
- Cao C, Lee X, Liu S, Schultz N, Xiao W, Zhang M and Zhao L 2016 Urban heat islands in China enhanced by haze pollution *Nat. Commun.* **7** 12509
- Cao J, Zhou W, Zheng Z, Ren T and Wang W 2021 Within-city spatial and temporal heterogeneity of air temperature and its relationship with land surface temperature *Landsc. Urban Plann.* **206** 103979
- Chakraborty T, Biswas T, Campbell L, Franklin B, Parker S and Tukman M 2022a Feasibility of afforestation as an equitable nature-based solution in urban areas *Sustain. Cities Soc.* **81** 103826
- Chakraborty T, Sarangi C and Tripathi S N 2017 Understanding diurnality and inter-seasonality of a sub-tropical urban heat island *Bound.-Layer Meteorol.* **163** 287–309
- Chakraborty T, Venter Z, Lee X and Qian Y 2022b Netatmo citizen weather station data over Europe (July, 2019) *Mendeley Data*, V1 (<https://doi.org/10.17632/ck69zx89fc.1>)
- Chakraborty T, Venter Z, Qian Y and Lee X 2022c Lower urban humidity moderates outdoor heat stress *AGU Adv.* **3** e2022AV000729
- Cheng X, Peng J, Dong J, Liu Y and Wang Y 2022 Non-linear effects of meteorological variables on cooling efficiency of African urban trees *Environ. Int.* **169** 107489
- Chinchilla J, Carbonnel A and Galleguillos M 2021 Effect of urban tree diversity and condition on surface temperature at the city block scale *Urban For. Urban Green.* **60** 127069
- Cleugh H 1998 Effects of windbreaks on airflow, microclimates and crop yields *Agrofor. Syst.* **41** 55–84
- Coutts A M, White E C, Tapper N J, Beringer J and Livesley S J 2016 Temperature and human thermal comfort effects of street trees across three contrasting street canyon environments *Theor. Appl. Clim.* **124** 55–68
- Dawson T E et al 2007 Nighttime transpiration in woody plants from contrasting ecosystems *Tree Physiol.* **27** 561–75
- Domec J-C et al 2012 Interactive effects of nocturnal transpiration and climate change on the root hydraulic redistribution and carbon and water budgets of southern United States pine plantations *Tree Physiol.* **32** 707–23
- Fan H, Yu Z, Yang G, Liu T Y, Liu T Y, Hung C H and Vejre H 2019 How to cool hot-humid (Asian) cities with urban trees? An optimal landscape size perspective *J. Agric. Meteorol.* **265** 338–48
- Goldblatt R, Addas A, Crull D, Maghrabi A, Levin G G and Rubinyi S 2021 Remotely sensed derived land surface temperature (LST) as a proxy for air temperature and thermal comfort at a small geographical scale *Land* **10** 410
- Grimm N B, Faeth S H, Golubiewski N E, Redman C L, Wu J, Bai X and Briggs J M 2008 Global change and the ecology of cities *Science* **319** 756–60
- Gui X, Wang L, Yao R, Yu D and Li C 2019 Investigating the urbanization process and its impact on vegetation change and urban heat island in Wuhan, China *Environ. Sci. Pollut. Res.* **26** 30808–25
- Gunawardena K R, Wells M J and Kershaw T 2017 Utilising green and bluespace to mitigate urban heat island intensity *Sci. Total Environ.* **584** 1040–55
- Haeri T, Hassan N and Ghaffarianhoseini A 2023 Evaluation of microclimate mitigation strategies in a heterogenous street canyon in Kuala Lumpur from outdoor thermal comfort perspective using Envi-met *Urban Clim.* **52** 101719
- Hami A, Abdi B, Zarehaghi D and Maulan S B 2019 Assessing the thermal comfort effects of green spaces: a systematic review of methods, parameters, and plants' attributes *Sustain. Cities Soc.* **49** 101634
- Harde H 2013 Radiation and heat transfer in the atmosphere: a comprehensive approach on a molecular basis *Int. J. Atmos. Sci.* **2013** 503727
- Hayes A T, Jandaghian Z, Lacasse M, Gaur A, Lu H, Laouadi A, Ge H and Wang L 2022 Nature-based solutions (nbss) to mitigate urban heat island (UHI) effects in Canadian cities *Buildings* **12** 925
- Ho H C, Knudby A, Xu Y, Hodul M and Aminipouri M 2016 A comparison of urban heat islands mapped using skin temperature, air temperature, and apparent temperature (Humidex), for the greater Vancouver area *Sci. Total Environ.* **544** 929–38
- Holzwarth S et al 2020 Earth observation based monitoring of forests in Germany: a review *Remote Sens.* **12** 3570
- Hsu A, Sheriff G, Chakraborty T and Manya D 2021 Disproportionate exposure to urban heat island intensity across major US cities *Nat. Commun.* **12** 2721
- Hulley G, Shivers S, Wetherley E and Cudd R 2019 New ECOSTRESS and MODIS land surface temperature data reveal fine-scale heat vulnerability in cities: a case study for Los Angeles County, California *Remote Sens.* **11** 2136
- Iungman T et al 2023 Cooling cities through urban green infrastructure: a health impact assessment of European cities *Lancet* **401** 577–89

- Janatian N, Sadeghi M, Sanaeinejad S H, Bakhshian E, Farid A, Hashemini S M and Ghazanfari S 2017 A statistical framework for estimating air temperature using MODIS land surface temperature data *Int. J. Climatol.* **37** 1181–94
- Jiao M, Zhou W, Zheng Z, Wang J and Qian Y 2017 Patch size of trees affects its cooling effectiveness: a perspective from shading and transpiration processes *J. Agric. Meteorol.* **247** 293–9
- Jin M S 2012 Developing an index to measure urban heat island effect using satellite land skin temperature and land cover observations *J. Clim.* **25** 6193–201
- Jin M and Dickinson R E 2010 Land surface skin temperature climatology: benefitting from the strengths of satellite observations *Environ. Res. Lett.* **5** 044004
- Kalnay E and Cai M 2003 Impact of urbanization and land-use change on climate *Nature* **423** 528–31
- Krayenhoff E S, Broadbent A M, Zhao L, Georgescu M, Middel A, Voogt J A, Martilli A, Sailor D J and Erell E 2021 Cooling hot cities: a systematic and critical review of the numerical modelling literature *Environ. Res. Lett.* **16** 053007
- Lagergren F and Lindroth A 2002 Transpiration response to soil moisture in pine and spruce trees in Sweden *J. Agric. Meteorol.* **112** 67–85
- Li H, Zhou Y, Jia G, Zhao K and Dong J 2022 Quantifying the response of surface urban heat island to urbanization using the annual temperature cycle model *Geosci. Front.* **13** 101141
- Li X, Chakraborty T C and Wang G 2023 Comparing land surface temperature and mean radiant temperature for urban heat mapping in Philadelphia *Urban Clim.* **51** 101615
- Li Y, Zhao M, Motesharrei S, Mu Q, Kalnay E and Li S 2015 Local cooling and warming effects of forests based on satellite observations *Nat. Commun.* **6** 6603
- Li Z-L, Wu H, Wang N, Qiu S, Sobrino J A, Wan Z, Tang B-H and Yan G 2013 Land surface emissivity retrieval from satellite data *Int. J. Remote Sens.* **34** 3084–127
- Liu H, Xu L, Wu J and Xu C 2019 Advances in studies on influential factors for cooling effect of urban forest *Sci. Silvae Sin.* **55** 144–51
- Liu Y, Huang X, Yang Q and Cao Y 2021 The turning point between urban vegetation and artificial surfaces for their competitive effect on land surface temperature *J. Clean. Prod.* **292** 126034
- Liu Y, Huang X, Yang Q, Jing W and Yang J 2023 Effects of landscape on thermal livability at the community scale based on fine-grained geographic information: a case study of Shenzhen *Sci. Total Environ.* **905** 167091
- Loughner C P, Allen D J, Zhang D-L, Pickering K E, Dickerson R R and Landry L 2012 Roles of urban tree canopy and buildings in urban heat island effects: parameterization and preliminary results *J. Appl. Meteorol. Climatol.* **51** 1775–93
- Lu Y, Lu S, Horton R and Ren T 2014 An empirical model for estimating soil thermal conductivity from texture, water content, and bulk density *Soil Sci. Soc. Am. J.* **78** 1859–68
- Meili N et al 2020 An urban ecohydrological model to quantify the effect of vegetation on urban climate and hydrology (UT&C v1.0) *Geosci. Model. Dev.* **13** 335–62
- Meili N et al 2021 Tree effects on urban microclimate: diurnal, seasonal, and climatic temperature differences explained by separating radiation, evapotranspiration, and roughness effects *Urban For. Urban Green.* **58** 126970
- Middel A, AlKhaled S, Schneider F A, Hagen B and Cosco P 2021 50 grades of shade *Bull. Am. Meteorol. Soc.* **102** E1805–20
- Mohajerani A, Bakaric J and Jeffrey-Bailey T 2017 The urban heat island effect, its causes, and mitigation, with reference to the thermal properties of asphalt concrete *J. Environ. Manage.* **197** 522–38
- Mutiibwa D, Strachan S and Albright T 2015 Land surface temperature and surface air temperature in complex terrain *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **8** 4762–74
- Naserikia M, Hart M A, Nazarian N, Bechtel B, Lipson M and Nice K A 2023 Land surface and air temperature dynamics: the role of urban form and seasonality *Sci. Total Environ.* **905** 167306
- Novick K A and Katul G G 2020 The duality of reforestation impacts on surface and air temperature *J. Geophys. Res. Biogeosci.* **125** e2019JG005543
- Pekel J-F, Cottam A, Gorelick N and Belward A S 2016 High-resolution mapping of global surface water and its long-term changes *Nature* **540** 418–22
- Peng -S-S, Piao S, Zeng Z, Ciais P, Zhou L, Li L Z X, Myneni R B, Yin Y and Zeng H 2014 Afforestation in China cools local land surface temperature *Proc. Natl Acad. Sci.* **111** 2915–9
- Peters E B, McFadden J P and Montgomery R A 2010 Biological and environmental controls on tree transpiration in a suburban landscape *J. Geophys. Res. Biogeosci.* **115** G04006
- Pirasteh-Anosheh H, Saed-Moucheshi A, Pakniyat H and Pesarakli M 2016 Stomatal responses to drought stress *Water Stress and Crop Plants: a Sustainable Approach* vol 1 pp 24–40
- Pramova E, Locatelli B, Djoudi H and Somorin O A 2012 Forests and trees for social adaptation to climate variability and change *Wiley Interdiscip. Rev. Clim. Change* **3** 581–96
- Rahman M A, Moser A, Rötzer T and Pauleit S 2017 Within canopy temperature differences and cooling ability of *Tilia cordata* trees grown in urban conditions *Build. Sci.* **114** 118–28
- Raman A P, Anoma M A, Zhu L, Rephaeli E and Fan S 2014 Passive radiative cooling below ambient air temperature under direct sunlight *Nature* **515** 540–4
- Ren J, Shi K, Li Z, Kong X and Zhou H 2023 A review on the impacts of urban heat islands on outdoor thermal comfort *Buildings* **13** 1368
- Richards D R, Fung T K, Belcher R and Edwards P J 2020 Differential air temperature cooling performance of urban vegetation types in the tropics *Urban For. Urban Green.* **50** 126651
- Schneider F A, Ortiz J C, Vanos J K, Sailor D J and Middel A 2023 Evidence-based guidance on reflective pavement for urban heat mitigation in Arizona *Nat. Commun.* **14** 1467
- Schwaab J, Meier R, Mussetti G, Seneviratne S, Bürgi C and Davin E L 2021 The role of urban trees in reducing land surface temperatures in European cities *Nat. Commun.* **12** 6763
- Tuholske C, Caylor K, Funk C, Verdin A, Sweeney S, Grace K, Peterson P and Evans T 2021 Global urban population exposure to extreme heat *Proc. Natl Acad. Sci.* **118** e2024792118
- Vanos J K, Middel A, Mc Kercher G R, Kuras E R and Ruddell B L 2016 Hot playgrounds and children's health: a multiscale analysis of surface temperatures in Arizona, USA *Landsch. Urban Plann.* **146** 29–42
- Venter Z S, Brousse O, Esau I and Meier F 2020 Hyperlocal mapping of urban air temperature using remote sensing and crowdsourced weather data *Remote Sens. Environ.* **242** 111791
- Venter Z S, Chakraborty T and Lee X 2021 Crowdsourced air temperatures contrast satellite measures of the urban heat island and its mechanisms *Sci. Adv.* **7** eabb9569
- Wan Z 2006 MODIS land surface temperature products users' guide *Institute for Computational Earth System Science (University of California)* p 805
- Wang C, Wang Z H and Yang J 2018 Cooling effect of urban trees on the built environment of contiguous United States *Earth's Future* **6** 1066–81
- Wang C, Wang Z-H, Wang C and Myint S W 2019 Environmental cooling provided by urban trees under extreme heat and cold waves in US cities *Remote Sens. Environ.* **227** 28–43
- Wang H, Wang X, Zhao P, Zheng H, Ren Y, Gao F and Ouyang Z 2012 Transpiration rates of urban trees, *Aesculus Chinensis* *J. Environ. Sci.* **24** 1278–87
- Wang J, Zhou W, Jiao M, Zheng Z, Ren T and Zhang Q 2020 Significant effects of ecological context on urban trees' cooling efficiency *ISPRS J. Photogramm. Remote Sens.* **159** 78–89
- Wang N, Chen J, He T, Xu X, Liu L, Sun Z, Qiao Z and Han D 2023a Understanding the differences in the effect of

- urbanization on land surface temperature and air temperature in China: insights from heatwave and non-heatwave conditions *Environ. Res. Lett.* **18** 104038
- Wang X, Dallimer M, Scott C E, Shi W and Gao J 2021 Tree species richness and diversity predicts the magnitude of urban heat island mitigation effects of greenspaces *Sci. Total Environ.* **770** 145211
- Wang X, Hsu A and Chakraborty T 2023b Citizen and machine learning-aided high-resolution mapping of urban heat exposure and stress *Environ. Res.:Infrastruct. Sustain.* **3** 035003
- Wujeska-Klause A and Pfautsch S 2020 The best urban trees for daytime cooling leave nights slightly warmer *Forests* **11** 945
- Yan C, Guo Q, Li H, Li L and Qiu G Y 2020 Quantifying the cooling effect of urban vegetation by mobile traverse method: a local-scale urban heat island study in a subtropical megacity *Build. Sci.* **169** 106541
- Yang Q, Huang X, Tong X, Xiao C, Yang J, Liu Y and Cao Y 2022 Global assessment of urban trees' cooling efficiency based on satellite observations *Environ. Res. Lett.* **17** 034029
- Yang Q, Xu Y, Tong X, Huang X, Liu Y, Chakraborty T C, Xiao C and Hu T 2023 An adaptive synchronous extraction (ASE) method for estimating intensity and footprint of surface urban heat islands: a case study of 254 North American cities *Remote Sens. Environ.* **297** 113777
- Yao R, Wang L, Huang X, Chen J, Li J and Niu Z 2018 Less sensitive of urban surface to climate variability than rural in Northern China *Sci. Total Environ.* **628** 650–60
- Zaitchik B F and Tuholske C 2021 Earth observations of extreme heat events: leveraging current capabilities to enhance heat research and action *Environ. Res. Lett.* **16** 111002
- Zeng Y et al 2023 Structural complexity biases vegetation greenness measures *Nat. Ecol. Evol.* **7** 1–9
- Zhang J, Gou Z, Zhang F and Shutter L 2020 A study of tree crown characteristics and their cooling effects in a subtropical city of Australia *Ecol. Eng.* **158** 106027
- Zhang M, Tan S, Zhang C, Han S, Zou S and Chen E 2023 Assessing the impact of fractional vegetation cover on urban thermal environment: a case study of Hangzhou, China *Sustain. Cities Soc.* **96** 104663
- Zhang P, Bounoua L, Imhoff M L, Wolfe R E and Thome K 2014 Comparison of MODIS land surface temperature and air temperature over the continental USA meteorological stations *Can. J. Remote Sens.* **40** 110–22
- Zhang T, Zhou Y, Zhu Z, Li X and Asrar G R 2022a A global seamless 1 km resolution daily land surface temperature dataset (2003–2020) *Earth Syst. Sci. Data* **14** 651–64
- Zhang Y, Wang Y, Ding N and Yang X 2022b Spatial pattern impact of impervious surface density on urban heat Island effect: a case study in xuzhou, China *Land* **11** 2135
- Zhao J, Zhao X, Wu D, Meili N and Fatichi S 2023a Satellite-based evidence highlights a considerable increase of urban tree cooling benefits from 2000 to 2015 *Glob. Change Biol.* **29** 3085–97
- Zhao Y, Li H, Bardhan R, Kubilay A, Li Q and Carmeliet J 2023b The time-evolving impact of tree size on nighttime street canyon microclimate: wind tunnel modeling of aerodynamic effects and heat removal *Urban Clim.* **49** 101528
- Zheng S, Guldmann J-M, Wang Z, Qiu Z, He C and Wang K 2021 Experimental and theoretical study of urban tree instantaneous and hourly transpiration rates and their cooling effect in hot and humid area *Sustain. Cities Soc.* **68** 102808
- Zhou D, Xiao J, Bonafoni S, Berger C, Deilami K, Zhou Y, Frolking S, Yao R, Qiao Z and Sobrino J 2018 Satellite remote sensing of surface urban heat Islands: progress, challenges, and perspectives *Remote Sens.* **11** 48
- Zhou S, L'Heureux M, Weaver S and Kumar A 2012 A composite study of the MJO influence on the surface air temperature and precipitation over the continental United States *Clim. Dyn.* **38** 1459–71
- Zhou W, Huang G, Pickett S T A, Wang J, Cadenasso M L, McPhearson T, Grove J M and Wang J 2021 Urban tree canopy has greater cooling effects in socially vulnerable communities in the US *One Earth* **4** 1764–75
- Zhou W, Wang J and Cadenasso M L 2017 Effects of the spatial configuration of trees on urban heat mitigation: a comparative study *Remote Sens. Environ.* **195** 1–12
- Ziter C D, Pedersen E J, Kucharik C J and Turner M G 2019 Scale-dependent interactions between tree canopy cover and impervious surfaces reduce daytime urban heat during summer *Proc. Natl Acad. Sci.* **116** 7575–80
- Zou Z et al 2021 Impacts of land use/land cover types on interactions between urban heat island effects and heat waves *Build. Sci.* **204** 108138